

# Fetal Organ Volume Measurements by Three-dimensional Ultrasonography in Clinical Practice

Rodrigo Ruano

## ABSTRACT

The present manuscript reviews the main three-dimensional ultrasound (3DUS) methods that are clinically available to measure fetal organ volumes. Nowadays, it is possible to measure the volume of different organs (lungs, heart, liver, kidneys, adrenal glands and brain) using 3DUS. Those measurements are clinically useful to predict pulmonary hypoplasia, cardiac dysfunction and anomalies, fetal growth, renal function and prematurity as well as to predict adequate and normal neuro-development.

**Keywords:** Fetal adrenal gland, Fetal brain volume, Fetal heart volume, Fetal kidney volume, Fetal liver volume, Fetal lung volume, Fetal organ volumes, Three-dimensional ultrasonography.

**How to cite this article:** Ruano R. Fetal Organ Volume Measurements by Three-dimensional Ultrasonography in Clinical Practice. *Donald School J Ultrasound Obstet Gynecol* 2015;9(4):397-407.

**Source of support:** Nil

**Conflict of interest:** None

## INTRODUCTION

The past three decades are remarkable for the impressive technological advance in ultrasound imaging methods. The three-dimensional ultrasonography (3DUS) represents one of these advances with the option to evaluate and measure fetal organ volumes in a three-dimensional image. Nowadays, new methods allow a volumetric measurement to be obtained faster and with more reproducibility. Particularly in maternal-fetal medicine, the possibility of evaluating the fetal organ volumes can impact in the perinatal management of the fetuses. In addition, new investigations have demonstrated that new methods can be used to estimate fetal weight in order to improve the diagnosis of intrauterine growth disorders. There are several studies that evaluated the impact of fetal

organ volumes estimated by 3DUS on the clinical prenatal management of those fetuses. This article presents the different methods of assessing volumes by 3DUS and its potential clinical applications in actual fetal medicine.

## DIFFERENT METHODS TO CALCULATE THE VOLUME USING THE THREE-DIMENSIONAL ULTRASONOGRAPHY

Basically, there are two simple and direct methods to assess the fetal organ volume using 3DUS approach.<sup>1</sup> The first one, which is the oldest method, consists in adding together the areas of serial parallel slices of the studied organ. The 'technique of the parallel slices' is based on 'cutting' the organ in many slices and measuring the areas from one side to another with specific pre-established thickness (Fig. 1). The main disadvantage of this method is that it may be difficult for small organs with irregular shapes.<sup>1</sup> Other limitations are described, such as the impossibility of repairing one incorrect area contouring.

More recently, a specific organ volume can be assessed by the rotational technique, which has become possible through the introduction of the 'virtual organ computer-aided analysis' (VOCAL<sup>TM</sup>) imaging program, an extension of the 3D View<sup>TM</sup> (GE, Kretztechnik, Zipf, Austria). This method allows measuring the organ volume by fixing one axis and then serial contouring the organ surface after rotating the volumetric image (Fig. 2). The volumetric image can be rotating using different angles. This is faster than the first method, and has the great advantage over the other in allowing corrections and verifications on the three-dimensional image. Despite of many studies have been demonstrating similar values, variations and reproducibility (agreement) with these two methods, nowadays the 'rotational technique' is more employed and preferable due to its facilities.<sup>1</sup>

## MEDICAL APPLICATIONS OF THE THREE-DIMENSIONAL VOLUMETRIC EVALUATION


### Fetal Weight Estimation

One of the main objectives of the prenatal care is the correct detection of fetal growth abnormalities. The large number of methods published in this respect is a clearly evidence of the importance of fetal weight estimation


Professor

Department of Obstetrics and Gynecology, Baylor College of Medicine, Houston, Texas, USA

**Corresponding Author:** Rodrigo Ruano, Professor, Department of Obstetrics and Gynecology, Baylor College of Medicine, 6651 Main Street, Suite F, Houston, Texas, USA, e-mail: rodrigoruan@hotmail.com



**Fig. 1:** Three-dimensional ultrasonographic parallel slices technique



**Fig. 2:** Three-dimensional ultrasonographic rotational technique rendered imaging of right and left fetal lung volumes

during prenatal care. Both fetal macrosomia and growth restriction increase perinatal morbidity and mortality. Fetuses with growth restriction are at increased risk of hypoxia and perinatal deaths. On the other hand, macrosomic fetuses are associated with increased risk of cesarean-section, postpartum hemorrhage, and maternal-fetal injuries. Macrosomic fetuses are generally associated with shoulder dystocia leading to clavicular fracture and brachial plexus injury.

Two-dimensional ultrasonography (2DUS) is applied routinely with this purpose. It estimates fetal weight indirectly by measuring defined segments of the body and then using specific mathematical equations, appropriate tables and integrated computer programs. The most frequently used fetal parameters are the biparietal diameter (BPD), head circumference (HC), abdominal circumference (AC) and femur length (Figs 3A to C).<sup>2-5</sup> There are so many formulas for estimating fetal weight, which is clear evidence that none of those is accepted universally. All these formulas for fetal weight estimation have a mean error of 7 to 10%, even in ideal ultrasound conditions.

The reason for the high error of these formulas may be the fact that none of these established equations takes the soft tissue thickness into account. It is known that fetal fat deposition represents approximately 90% of caloric accretion at term, and that neonatal fat mass constitutes 12 to 15% of birth weight. There have been many studies



**Figs 3A to C:** Fetal parameters for fetal weight estimation: (A) Biparietal diameter and head circumference, (B) abdominal circumference and (C) femur length

using different methods to take the soft-tissue content into account on 2DUS, especially by including the limb diameters, with varying results.<sup>6</sup>

Recently, with the advent of new volumetric methods using 3DUS technology, measurements of circumference and volume have become possible and reproducible through the three orthogonal sections of the specific fetal organ. Regarding the potential use for the estimation of fetal weight, the main interest becomes the possibility of measuring fetal limb diameters and volumes on these sections, which has the advantages of being more efficient and faster than the 2DUS technologies.<sup>6-9</sup> In addition, these new methods allow incorporation of direct measurements of the fetal soft tissue in the equations to estimate fetal weight. The method used to measure the fetal limb volumes is called fractional limb volumes.<sup>6-9</sup>

The fractional limb volume is a method that the fetal soft tissue is evaluated from a central portion of the limb (arm and thigh) diaphysis (Figs 4 and 5). Transverse slices of the mid-limb are more likely to display the soft tissue borders. This technique reduces the measuring times and also improves the reproducibility of the method. Tables 1 and 2 present different mathematical equations and reference values of the limb volumes by gestational age.

### Fetal Organ Volume

Three-dimensional ultrasound was first introduced during the 1980s, and has gained increasing acceptance in obstetrics as technological improvement. Its use has revolutionized the prenatal detection of fetal structural anomalies, growth aberrations, and multiple gestations. In recent years, 3DUS has undergone rapid advances, and continues to progress quickly. More recently, the estimation of fetal organ volumes is a promising application in maternal-fetal medicine. Different studies have




Fig. 4: Fractional thigh volume




Fig. 5: Fractional arm volume

proposed the use of 3DUS to measure different fetal organ volumes (heart, liver, lung, brain, renal, adrenal gland and thymus). Unquestionably, 3DUS of fetal organs is gaining increasing importance in prenatal diagnosis, and will likely become an integral part of it in the near future.

The general essential prerequisites for measurement of a fetal organ volume are: first, acquisition of the volume with 'the organ of interest' facing the transducer; second, this acquisition must be in the absence of fetal

movements and, third, accurate definition of the whole organ with good identification of its borders. Gestational age may interfere with the quality of the volumetric acquisition, and may limit posterior analysis of the 'organ of interest'. For example, lung volumes are more reproducible before 32 weeks, differently of the kidney volumes. Other important point is that acquisition and volumetric analysis depend on the amount of amniotic fluid and other organs shadows.



**Table 1:** Fetal weight estimation formulae using 3DUS measurements

Author	Regression equation
Lee et al 2001 <sup>6</sup>	$FW = 20.95 \times Tvol + 113.571 \times AC - 2375.068$
Lee et al 2009 <sup>7</sup>	$\text{LogFW} = 11.1372 \times \text{BPD}^2 - 67.2281 \times \text{BPD} + 1.2175 \times \text{AC}^2 - 17.3004 \times \text{AC} - 0.0490 \times \text{Tvol}^2 + 25.3052 \times \text{Tvol} + 285.429$

FW: Fetal weight; BPD: Biparietal diameter; HC: Head circumference; AC: Abdominal circumference; FL: Femur length; Tvol: Thigh volume; ABDvol: Abdominal volume

**Table 2:** Percentiles of fetal humerus volume (cm<sup>3</sup>) estimated by 3DUS using the rotational technique (VOCAL™) in accordance to gestational age in weeks

GA	Fetal humerus volume (cm <sup>3</sup> )		
	Percentiles		
	10%	50%	90%
20	0.71	0.77	0.82
22	0.86	0.97	1.07
24	1.05	1.20	1.35
26	1.28	1.48	1.67
28	1.54	1.78	2.02
30	1.84	2.12	2.41
32	2.17	2.50	2.83
34	2.54	2.91	3.29
36	2.94	3.36	3.79
38	3.38	3.85	4.32
40	3.85	4.36	4.88

Modified from Chang et al. *Ultrasound in Medicine and Biology* 2005<sup>9</sup>

## Fetal Lung Volume

Clinically, fetal lung volume measurements can be useful to predict the severity of postnatal pulmonary hypoplasia more accurately in determined conditions, such as congenital diaphragmatic hernia, hyperechogenic lung lesions and oligohydramnios. Conventional 2DUS, which is used routinely in obstetrics, has proved to be insufficiently reliable to diagnose and predict prognosis of determinate congenital malformation, e.g. in fetuses at high risk for pulmonary hypoplasia. Recent studies have demonstrated the use of 3DUS in predicting pulmonary hypoplasia.<sup>10-21</sup> This technology has the same advantages as 2DUS related to cost-effectiveness, feasible and speed of use and patient acceptability, but in addition, enables visualization of perpendicular planes simultaneously (multiplanar imaging) and acquisition of the entire organ volume.

Nowadays, studies demonstrate that both methods of measuring fetal lung volumes can be used, the parallel slices or the rotational technique. However, the rotational technique is preferable for small and irregular lungs, especially to evaluate lung volumes in congenital diaphragmatic hernia with better resolution (Figs 6A to D).

Assessing fetal lung volumes have been shown to be one of the best ways to predict neonatal prognosis in cases with congenital diaphragmatic hernia.<sup>10,12,14,18-21</sup> The importance of predicting neonatal outcome in those fetuses is based on the possibility to select cases for antenatal interventions. Actually, fetuses with congenital diaphragmatic hernia presenting worse prognosis with high risk to develop severe pulmonary hypoplasia after birth may benefit from transitory tracheal occlusion during fetal life.<sup>22-30</sup> The advantages of evaluating fetal lung volumes by 3DUS over assessing measurements on 2DUS are that the entire lung is considered in one image alone and that the lung ipsilateral to the diaphragmatic defect may be included in the analysis.

Predicting the severity of pulmonary hypoplasia may also be useful in fetuses with large hyperechogenic pulmonary lesions (adenomatoid malformation or pulmonary sequestration). For those fetuses, depending on the severity of the pulmonary hypoplasia, a few cases will benefit for early postnatal intervention such *ex utero* intrapartum treatment (EXIT) and extracorporeal membrane oxygenation (ECMO).<sup>31-33</sup>


More recent studies have also suggested that fetuses with very early and prolonged premature rupture of the membranes may also benefit from fetal tracheal occlusion.<sup>34-36</sup> Fetal lung volumetric analysis may also be useful for the selection process in those cases.

There are so many reports on normal values and ranges of fetal lung volumes throughout gestational age. We have demonstrated that the right, left and total lung volumes vary from 5.37, 4.66 and 9.95 cm<sup>3</sup> at 20 weeks to 46.6, 37.34, and 84.35 cm<sup>3</sup> at 37 weeks, based on mathematical formulas: right lung volume =  $\exp \{4.07/[1 + \exp (21.90 - \text{gestational age}/5.44)]\}$ ; left lung volume =  $\exp \{3.82/[1 + \exp (22.03 - \text{gestational age}/5.17)]\}$ , and total lung volume =  $\exp \{4.72/[1 + \exp (20.30 - \text{gestational age}/6.05)]\}$  (Table 3).<sup>15</sup>

## Fetal Heart Volume

Congenital heart disease is a leading cause of infant mortality in many countries, with an estimated incidence of about 3 to 8 per 1000 live births. Prenatal diagnosis of congenital heart diseases is responsible for a better planning of the perinatal management improving the postnatal care and outcomes.

Heart volume is a valuable parameter used in the evaluation of fetal heart condition, being mandatory for early detection of impending collapse of fetal circulation. Its accurate assessment becomes indispensable for prenatal diagnosis and management of fetal cardiac situations, such as in fetal high-output failure, hydrops fetalis, and many clinical-pathological conditions. The classical



**Figs 6A to D:** Fetal lung volume: (A) Transverse section, (B) sagittal section, (C) coronal section and (D) rendered imaging

**Table 3:** Percentiles of fetal lung volume ( $\text{cm}^3$ ) estimated by 3DUS using the rotational technique (VOCAL™) in accordance to gestational age in weeks

GA	Right fetal lung volume ( $\text{cm}^3$ )			Left fetal lung volume ( $\text{cm}^3$ )			Total fetal lung volume ( $\text{cm}^3$ )		
	Percentiles			Percentiles			Percentiles		
	10%	50%	90%	10%	50%	90%	10%	50%	90%
20	4.31	5.37	6.75	3.60	4.66	6.05	8.03	9.95	12.45
22	6.23	7.77	9.68	5.21	6.69	8.58	11.59	14.66	18.28
24	8.94	11.25	14.15	7.61	9.68	12.30	16.83	21.22	26.88
26	12.81	15.96	20.09	10.28	13.60	17.12	23.66	29.70	37.52
28	17.12	21.54	27.11	14.15	18.36	23.34	31.97	39.74	50.17
30	21.98	27.66	34.47	18.73	23.34	29.96	40.73	50.63	63.99
32	27.39	33.78	42.95	21.98	28.22	35.87	49.47	61.50	76.90
34	32.14	39.25	49.90	24.78	32.46	41.68	57.84	71.62	92.03
36	35.87	43.82	54.60	27.66	35.87	46.53	64.55	80.47	101.85

Modified from Ruano, et al. Journal of Ultrasound in Medicine 2006<sup>15</sup>

method using 2DUS for assessing heart volume, which assumes that heart shape is spherical or elliptical is imprecise. However, before the advent of an appropriate noninvasive research tool, such as 3DUS, this was difficult to be evaluated in human fetuses.


Traditionally, fetal heart volume has been assessed by the equation: heart volume =  $0.5233 \times H1 \times H2 \times H2$ , where H1 is the largest longitudinal diameter of the fetal heart at the four-chamber view, and H2 is the largest transverse diameter at the same plane.

Fetal heart volumes can be evaluated by the rotational technique on 3DUS (Figs 7A to D).<sup>37,38</sup> There have been a very few reports on normal values of fetal heart volume throughout gestational age. Fetal heart volume may vary

from  $0.63$  ( $0.40$ – $0.86$ )  $\text{cm}^3$  at 12 weeks to  $26.60$  ( $16.40$ – $36.80$ )  $\text{cm}^3$  at 32 weeks (Table 4).<sup>37,39</sup>

### Fetal Liver Volume

Determination of fetal liver volume is very useful in assessing the status of fetal growth and nutrition. Fetuses with intrauterine growth restriction or macrosomia have different liver volumes. However, to measure human fetal liver *in utero* precisely and noninvasively is not an easy task. In the past, many studies have performed several attempts to indirectly assess liver size, using different parameters, such as AC and diameters (anteroposterior, transverse and cephalocaudal) using 2DUS, and then multiplied by a constant which had been obtained



**Figs 7A to D:** Fetal heart volume: (A) Transverse section, (B) sagittal section, (C) coronal section and (D) rendered imaging


from adult liver studies, to estimate fetal liver weight. Currently, fetal liver volume is measured using its longest diameter, which may be less reproducible. With the advent of 3DUS and the improvement of its technology, fetal liver volume assessment becomes possible, easier and more reproducible (Figs 8A to D).<sup>40-47</sup> Besides, it has become an easy and noninvasive approach to obtain precise measurement of fetal liver volumes, and good correlation has been established between liver volume and other fetal growth parameters.

There have been very few studies reporting on normal ranges of fetal liver volumes assessed by 3DUS. Fetal liver

volumes range from 11.73 cm<sup>3</sup> at 20 weeks to 131.51 cm<sup>3</sup> at 40 weeks (Table 5).<sup>42,47</sup>

### Fetal Renal Volume

Renal diseases are not uncommon in fetal life, and some of them might be treatable or carry a better outcome if detected earlier. Evaluating fetal renal size may be a valuable tool in the detection and monitoring of renal conditions, including nephromegaly, hypoplasia and other anomalies. Besides, assessing fetal renal size may be useful to evaluate renal function. Konje et al<sup>48</sup> demonstrated that fetal renin concentration is related to



**Figs 8A to D:** Fetal liver volume: (A) Transverse section, (B) sagittal section, (C) coronal section and (D) rendered imaging

**Table 4:** Percentiles of fetal heart volume (cm<sup>3</sup>) estimated by 3DUS using the rotational technique (VOCAL™) in accordance to gestational age in weeks

GA	Fetal heart volume (cm <sup>3</sup> )		
	Percentiles		
	2.50%	50%	97.50%
12	0.40	0.63	0.86
14	0.17	0.53	0.88
16	0.46	1.14	1.81
18	1.24	2.42	3.61
20	2.44	4.33	6.23
22	4.04	6.83	9.62
24	5.99	9.87	13.80
26	8.24	13.40	18.60
28	10.80	17.40	24.00
30	13.50	21.80	30.10
32	16.40	26.60	36.80

Modified from Peralta, et al. *Ultrasound in Obstetrics and Gynecology* 2006<sup>39</sup>

**Table 5:** Percentiles of fetal liver volume (cm<sup>3</sup>) estimated by 3DUS using the rotational technique (VOCAL™) in accordance to gestational age in weeks

GA	Fetal liver volume (cm <sup>3</sup> )		
	Percentiles		
	10%	50%	90%
20	5.05	11.84	18.63
22	11.82	19.17	26.51
24	17.00	24.90	32.80
26	21.73	30.18	38.63
28	27.18	36.13	45.13
30	34.32	43.38	53.44
32	44.46	54.57	64.69
34	58.67	69.34	80.00
36	78.08	83.30	100.53
38	103.83	115.61	127.38
40	137.05	149.38	161.71

Modified from Chang et al. *Ultrasound in Medicine and Biology* 2003<sup>42</sup>


fetal size. The classical way of measuring fetal renal size using 2DUS is not precise. With the application of 3DUS, it becomes feasible and more accurate by estimating fetal renal volumes (Figs 9A to D).

There have been very few reports on the distribution of fetal renal volumes throughout gestational age. Hsieh et al<sup>49</sup> described good correlation between fetal renal volumes and gestational age, suggesting the following mathematical formulas: right fetal renal volume =  $0.4032 \times \text{gestational age} - 6.2097$ ; left fetal renal volume =  $0.3954 \times \text{gestational age} - 6.3068$ . Yu et al<sup>50</sup> reported the following mathematical formulas: right fetal renal volume =  $0.74053 \times \text{gestational week} - 13.318$ ; left fetal renal volume =  $0.76093 \times \text{gestational age} - 13.421$ .

More recently, Yoshizaki et al<sup>51</sup> reported the following mathematical formulas: expected right kidney volume =  $\exp[-1.01 + (0.12 \times \text{gestational age})]$ , and expected left kidney volume =  $\exp[-0.90 + (0.12 \times \text{gestational age})]$  (Table 6).

**Fetal Adrenal Gland Volume**

The fetal adrenal glands are important organs to fetal growth. *In utero*, the normal fetal adrenal glands can be scanned using 2DUS as hypoechoic structures located immediately above the fetal kidneys bilaterally. Some pathologic conditions of fetal adrenal glands, such as adrenoblastomas or neuroblastomas, can be diagnosed on ultrasound examinations. These fetal adrenal tumors



**Figs 9A to D:** Fetal renal volume: (A) Sagittal section, (B) transverse section, (C) coronal section and (D) rendered imaging



**Table 6:** Percentiles of fetal renal volume (cm<sup>3</sup>) estimated by 3DUS using the rotational technique (VOCAL™) in accordance to gestational age in weeks

GA	Right fetal renal volume (cm <sup>3</sup> )			Left fetal renal volume (cm <sup>3</sup> )		
	Percentiles			Percentiles		
	10%	50%	90%	10%	50%	90%
20	2.4	4.0	6.9	2.4	4.2	7.4
22	3.0	5.1	8.7	3.1	5.4	9.4
24	3.9	6.5	11.1	3.9	6.8	11.8
26	4.9	8.3	14.1	4.9	8.6	15.0
28	6.3	10.6	18.0	6.2	10.8	18.9
30	8.0	13.5	22.9	7.9	13.7	23.9
32	10.1	17.2	29.1	9.9	17.3	30.2
34	12.9	21.8	37.0	12.6	21.9	38.2
36	14.1	27.8	47.1	15.9	27.7	48.4
38	20.8	35.3	59.9	20.1	35.1	61.2
40	26.5	45.0	76.2	25.4	44.3	77.3

Modified from Yoshizaki, et al. J Ultrasound Med 2013<sup>51</sup>

may have abnormal sizes. Therefore, measuring this gland volume may be useful for the early diagnosis of this condition.

The irregular shape of fetal adrenal glands makes difficult to evaluate its volume using 2DUS. With the advent of 3DUS, using the rotational technique, it becomes feasible and noninvasive method to evaluate its volumes (Figs 10A to D).<sup>52-57</sup> Reference values may vary according to the following equation: adrenal gland volume =  $-0.2683 \times \text{gestational age} + 0.0082 \times (\text{gestational age})^2 + 3.1927$  (Table 7).<sup>52</sup>

Other interest in assessing fetal adrenal gland volumes can be related to a potential use in predicting labor, especially preterm labor.<sup>53,55</sup> Several mechanisms have

**Table 7:** Percentiles of fetal adrenal gland volume (cm<sup>3</sup>) estimated by 3DUS using the rotational technique (VOCAL™) in accordance to gestational age in weeks


GA	Fetal adrenal gland volume (cm <sup>3</sup> )		
	Percentiles		
	10%	50%	90%
22	0.86	1.24	1.63
24	1.01	1.45	1.90
26	1.22	1.73	2.25
28	1.50	2.08	2.65
30	1.85	2.49	3.13
32	2.26	2.96	3.67
34	2.74	3.51	4.27
36	3.28	4.11	4.94
38	3.89	4.78	5.68
40	4.56	5.52	6.48

Modified from Chang et al. Ultrasound in Medicine and Biology 2002<sup>52</sup>

been implicated in triggering preterm labor, like genetic predisposition, decidual hemorrhage, fetal stress and inflammation/infections. In most animal species, the fetal hypothalamic-pituitary-adrenal axis plays a special role in the initiation of parturition.

**Fetal Brain Volume**


Both fetal BPD and fetal HC are standard parameters in establishing normal and abnormal fetal biometry and growth. Fetal brain volume measurement in conjunction with fetal liver volume determination could provide insight into the nature of abnormal fetal growth.<sup>43</sup> Up-to-date, there have been few studies reporting on the use of 3DUS to assess fetal brain volume (Figs 11A to D). Fetal brain volumes vary from 53 cm<sup>3</sup> at 20 weeks to 316 cm<sup>3</sup>



**Figs 10A to D:** Fetal adrenal gland volume: (A) Sagittal section, (B) transverse section, (C) coronal section and (D) rendered imaging







**Figs 11A to D:** Fetal brain volume: (A) Transverse section, (B) sagittal section, (C) coronal section and (D) rendered imaging

**Table 8:** Percentiles of fetal brain volume ( $\text{cm}^3$ ) estimated by 3DUS using the rotational technique (VOCAL<sup>TM</sup>) in accordance to gestational age in weeks

GA	Fetal brain volume ( $\text{cm}^3$ )		
	Percentiles		
	5%	50%	95%
20	38	53	71
22	58	77	100
24	82	116	133
26	109	139	173
28	140	177	218
30	174	219	269
32	211	265	326
34	252	316	389

Modified from Roelfsema et al. American Journal of Obstetrics and Gynecology 2004<sup>58</sup>

at 34 weeks according to the following equation: fetal brain volume =  $0.75 \times \text{gestational age} - 7.71$  (Table 8).<sup>58</sup>

In addition to the entire fetal brain volume, it is possible to measure the fetal hippocampus, cerebellum, thalamus and ventricles.<sup>59-66</sup> Recent studies have demonstrated that fetuses with particular diseases, such as congenital heart diseases have not only smaller brain, but reduced volumes of intracranial structures.<sup>67</sup> Three-dimensional ultrasound is an important prenatal diagnostic tool in fetal neurosonography, which can be used to predict adequate neurodevelopment.<sup>68,69</sup>

## CONCLUSION

Nowadays, it is possible to evaluate the fetal organ volumes using 3DUS technology. The main advantage of using 3DUS over 2DUS technology in measuring the organ volume is related to the fact that the first method

permits evaluation of the entire organ instead of part of it. Using the rotational technique, the process of measuring the fetal organ volumes has become more feasible and reproducible. There is still a need to standardize and generalize the methods of measuring the fetal organ volumes. However, several clinical applications of 3DUS fetal organ measurements have been demonstrated.

## REFERENCES

1. Ruano R. Recent advances in sonographic imaging of fetal thoracic structures. *Expert Review of Medical Devices* 2005;2(2):217-222.
2. Hadlock FP, Harrist RB, Martinez-Poyer J. In utero analysis of fetal growth: a sonographic weight standard. *Radiology* 1991;181(1):129-133.
3. Deter RL, Rossavik IK, Harrist RB, Hadlock FP. Mathematic modeling of fetal growth: development of individual growth curve standards. *Obstet Gynecol* 1986;68(2):156-161.
4. Hadlock FP, Deter RL, Harrist RB, Roecker E, Park SK. A date-independent predictor of intrauterine growth retardation: femur length/abdominal circumference ratio. *Am J Roentgenol* 1983;141(5):979-984.
5. Deter RL, Harrist RB, Hadlock FP, Poindexter AN. Longitudinal studies of fetal growth with the use of dynamic image ultrasonography. *Am J Obstet Gynecol* 1982;143(5):545-554.
6. Lee W, Deter RL, Ebersole JD, Huang R, Blanckaert K, Romero R. Birth weight prediction by three-dimensional ultrasonography: fractional limb volume. *J Ultrasound Med.* 2001;20(12):1283-1292.
7. Lee W, Balasubramaniam M, Deter RL, Yeo L, Hassan SS, Gotsch F, Kusanovic JP, Goncalves LF, Romero R. New fetal weight estimation models using fractional limb volume. *Ultrasound Obstet Gynecol* 2009;34(5):556-565.
8. Lee W, Deter R, Sangi-Haghpeykar H, Yeo L, Romero R. Prospective validation of fetal weight estimation using fractional limb volume. *Ultrasound Obstet Gynecol* 2013;41(2):198-203.

9. Chang CH, Yu CH, Ko HC, Chen CL, Chang FM. Fetal upper arm volume in predicting intrauterine growth restriction: a three-dimensional ultrasound study. *Ultrasound Med Biol* 2005;31(11):1435-1439.
10. Ruano R, Benachi A, Martinovic J, Grebille AG, Aubry MC, Dumez Y, Dommergues M. Can three-dimensional ultrasound be used for the assessment of the fetal lung volume in cases of congenital diaphragmatic hernia? *Fetal Diagnosis and Therapy* 2004;19(1):87-91.
11. Sabogal JC, Becker E, Bega G, Komwilaisak R, Berghella V, Weiner S, Tolosa J. Reproducibility of fetal lung volume measurements with 3-dimensional ultrasonography. *J Ultrasound Med* 2004;23(3):347-352.
12. Ruano R, Martinovic J, Dommergues M, Aubry MC, Dumez Y, Benachi A. Accuracy of fetal lung volume assessed by three-dimensional sonography. *Ultrasound Obstet Gynecol* 2005;26(7):725-730.
13. Gerards FA, Engels MA, Twisk JW, van Vugt JM. Normal fetal lung volume measured with three-dimensional ultrasound. *Ultrasound Obstet Gynecol*. 2006;27(2):134-144.
14. Ruano R, Aubry MC, Barthe B, Mitanchez D, Dumez Y, Benachi A. Quantitative analysis of fetal pulmonary vasculature by 3-dimensional power Doppler ultrasonography in isolated congenital diaphragmatic hernia. *Am J Obstetrics Gynecol* 2006;195(6):1720-1728.
15. Ruano R, Joubin L, Aubry MC, Thalabard JC, Dommergues M, Dumez Y, Benachi A. A nomogram of fetal lung volumes estimated by 3-dimensional ultrasonography using the rotational technique (virtual organ computer-aided analysis). *Ultrasound Med* 2006;25(6):701-709.
16. Gerards FA, Twisk JW, Bakker M, Barkhof F, van Vugt JM. Fetal lung volume: three-dimensional ultrasonography compared with magnetic resonance imaging. *Ultrasound Obstet Gynecol* 2007;29(5):533-536.
17. Hata T, Kuno A, Dai SY, Inubashiri E, Hanaoka U, Kanenishi K, Yamashiro C, Tanaka H, Yanagihara T. Three-dimensional sonographic volume measurement of the fetal lung. *J Obstet Gynaecol Res* 2007;33(6):793-798.
18. Ruano R, Aubry MC, Barthe B, Dumez Y, Zugaib M, Benachi A. Ipsilateral lung volumes assessed by three-dimensional ultrasonography in fetuses with isolated congenital diaphragmatic hernia. *Fetal Diagnosis and Therapy* 2008;24(4):389-394.
19. Kehl S, Eckert S, Sutterlin M, Neff KW, Siemer J. Influence of different rotation angles in assessment of lung volumes by 3-dimensional sonography in comparison to magnetic resonance imaging in healthy fetuses. *J Ultrasound Med* 2011;30(6):819-825.
20. Prendergast M, Rafferty GF, Davenport M, Persico N, Jani J, Nicolaides K, Greenough A. Three-dimensional ultrasound fetal lung volumes and infant respiratory outcome: a prospective observational study. *BJOG: an International Journal of Obstetrics and Gynaecology* 2011;118(5):608-614.
21. Kehl S, Becker L, Eckert S, Weiss C, Schaible T, Neff KW, Siemer J, Sutterlin M. Prediction of mortality and the need for neonatal extracorporeal membrane oxygenation therapy by 3-dimensional sonography and magnetic resonance imaging in fetuses with congenital diaphragmatic hernias. *J Ultrasound Med* 2013;32(6):981-988.
22. Deprest J, Gratacos E, Nicolaides KH. Fetoscopic tracheal occlusion (FETO) for severe congenital diaphragmatic hernia: evolution of a technique and preliminary results. *Ultrasound Obstet Gynecol* 2004;24(2):121-126.
23. Deprest JA, Hyett JA, Flake AW, Nicolaides K, Gratacos E. Current controversies in prenatal diagnosis 4: Should fetal surgery be done in all cases of severe diaphragmatic hernia? *Prenat Diagn* 2009;29(1):15-19.
24. Deprest JA, Nicolaides K, Gratacos E. Fetal surgery for congenital diaphragmatic hernia is back from never gone. *Fetal Diagn Ther* 2011;29(1):6-17.
25. Ruano R, Duarte SA, Pimenta EJ, Takashi E, da Silva MM, Tannuri U, Zugaib M. Comparison between fetal endoscopic tracheal occlusion using a 1.0 mm fetoscope and prenatal expectant management in severe congenital diaphragmatic hernia. *Fetal Diagn Ther* 2011;29(1):64-70.
26. Deprest J, De Coppi P. Antenatal management of isolated congenital diaphragmatic hernia today and tomorrow: ongoing collaborative research and development. *J Pediatric Surg Lecture. J Pediatr Surg* 2012;47(2):282-290.
27. Ruano R, da Silva MM, Campos JA, Papanna R, Moise K, Jr, Tannuri U, Zugaib M. Fetal pulmonary response after fetoscopic tracheal occlusion for severe isolated congenital diaphragmatic hernia. *Obstet gynecol* 2012;119(1):93-101.
28. Ruano R, Yoshisaki CT, da Silva MM, Ceccon ME, Grasi MS, Tannuri U, Zugaib M. A randomized controlled trial of fetal endoscopic tracheal occlusion versus postnatal management of severe isolated congenital diaphragmatic hernia. *Ultrasound Obstet Gynecol* 2012;39(1):20-27.
29. Done E, Gratacos E, Nicolaides KH, Allegaert K, Valencia C, Castanon M, Martinez JM, Jani J, Van Mieghem T, Greenough A, et al. Predictors of neonatal morbidity in fetuses with severe isolated congenital diaphragmatic hernia undergoing fetoscopic tracheal occlusion. *Ultrasound Obstet and Gynecol* 2013;42(1):77-83.
30. Ruano R, Peiro JL, da Silva MM, Campos JA, Carreras E, Tannuri U, Zugaib M. Early fetoscopic tracheal occlusion for extremely severe pulmonary hypoplasia in human isolated congenital diaphragmatic hernia: preliminary results. *Ultrasound Obstet Gynecol* 2013.
31. Keller TM, Rake A, Michel SC, Seifert B, Wissler J, Marincek B, Kubik-Huch RA. MR assessment of fetal lung development using lung volumes and signal intensities. *Eur Radiol* 2004;14(6):984-989.
32. Matsushita M, Ishii K, Tamura M, Takahashi Y, Kamura T, Takakuwa K, Tanaka K. Perinatal magnetic resonance fetal lung volumetry and fetal lung-to-liver signal intensity ratio for predicting short outcome in isolated congenital diaphragmatic hernia and cystic adenomatoid malformation of the lung. *J Obstet Gynaecol Res* 2008;34(2):162-167.
33. Rypens F, Metens T, Rocourt N, Sonigo P, Brunelle F, Quere MP, Guibaud L, Maugey-Laulom B, Durand C, Avni FE, et al. Fetal lung volume: estimation at MR imaging-initial results. *Radiology* 2001;219(1):236-241.
34. Kohl T, Geipel A, Tchatcheva K, Stressig R, Willinek WA, Gembruch U, Muller A. Life-saving effects of fetal tracheal occlusion on pulmonary hypoplasia from preterm premature rupture of membranes. *Obstet Gynecol* 2009;113(2 Pt 2):480-483.
35. Barth WH, Jr. Fetoscopic tracheal occlusion for previsible rupture of the membranes: recklessness or heroic medical innovation? *Obstet Gynecol* 2009;113(2 Pt 2):473-475.
36. Kohl T, Muller A, Franz A, Heep A, Willinek WA, Bartmann P, Gembruch U. Temporary fetoscopic tracheal balloon occlusion enhanced by hyperoncotic lung distension: is there a role in the treatment of fetal pulmonary hypoplasia from early preterm premature rupture of membranes? *Fetal Diagn Ther* 2007;22(6):462-465.
37. Chang FM, Hsu KF, Ko HC, Yao BL, Chang CH, Yu CH, Liang RI, Chen HY. Fetal heart volume assessment by three-dimensional ultrasound. *Ultrasound Obstet Gynecol* 1997;9(1):42-48.
38. Herberg U, Luck S, Steinweg B, Brand M, Knies R, Geipel A, Trier HG, Breuer J. Volumetry of fetal hearts using 3D real-



- time matrix echocardiography-in vitro validation experiments and 3D echocardiographic studies in fetuses. *Ultraschall Med* 2011;32(1):46-53.
39. Peralta CF, Cavoretto P, Csapo B, Falcon O, Nicolaides KH. Lung and heart volumes by three-dimensional ultrasound in normal fetuses at 12-32 weeks gestation. *Ultrasound Obstet Gynecol* 2006;27(2):128-133.
  40. Chang CH, Yu CH, Ko HC, Chen CL, Chang FM. Predicting fetal growth restriction with liver volume by three-dimensional ultrasound: efficacy evaluation. *Ultrasound Med Biol* 2006;32(1):13-17.
  41. Boito SM, Struijk PC, Ursem NT, Stijnen T, Wladimiroff JW. Assessment of fetal liver volume and umbilical venous volume flow in pregnancies complicated by insulin-dependent diabetes mellitus. *BJOG: an International Journal of Obstetrics and Gynaecology* 2003;110(11):1007-1013.
  42. Chang CH, Yu CH, Chang FM, Ko HC, Chen HY. The assessment of normal fetal liver volume by three-dimensional ultrasound. *Ultrasound Med Biol* 2003;29(8):1123-1129.
  43. Boito S, Struijk PC, Ursem NT, Fedele L, Wladimiroff JW. Fetal brain/liver volume ratio and umbilical volume flow parameters relative to normal and abnormal human development. *Ultrasound Obstet Gynecol* 2003;21(3):256-261.
  44. Boito SM, Laudy JA, Struijk PC, Stijnen T, Wladimiroff JW. Three-dimensional US assessment of hepatic volume, head circumference, and abdominal circumference in healthy and growth-restricted fetuses. *Radiology* 2002;223(3):661-665.
  45. Kuno A, Hayashi Y, Akiyama M, Yamashiro C, Tanaka H, Yanagihara T, Hata T. Three-dimensional sonographic measurement of liver volume in the small-for-gestational-age fetus. *J Ultrasound Med* 2002;21(4):361-366.
  46. Laudy JA, Janssen MM, Struyk PC, Stijnen T, Wallenburg HC, Wladimiroff JW. Fetal liver volume measurement by three-dimensional ultrasonography: a preliminary study. *Ultrasound Obstet Gynecol* 1998;12(2):93-96.
  47. Chang FM, Hsu KF, Ko HC, Yao BL, Chang CH, Yu CH, Chen HY. Three-dimensional ultrasound assessment of fetal liver volume in normal pregnancy: a comparison of reproducibility with two-dimensional ultrasound and a search for a volume constant. *Ultrasound Med Biol* 1997;23(3):381-389.
  48. Konje JC, Bell SC, Morton JJ, de Chazal R, Taylor DJ. Human fetal kidney morphometry during gestation and the relationship between weight, kidney morphometry and plasma active renin concentration at birth. *Clin Sci (Lond)* 1996;91(2):169-175.
  49. Hsieh YY, Chang CC, Lee CC, Tsai HD. Fetal renal volume assessment by three-dimensional ultrasonography. *Am J Obstet Gynecol* 2000;182(2):377-379.
  50. Yu C, Chang C, Chang F, Ko H, Chen H. Fetal renal volume in normal gestation: a three-dimensional ultrasound study. *Ultrasound Med Biol* 2000;26(8):1253-1256.
  51. Yoshizaki CT, Francisco RP, de Pinho JC, Ruano R, Zugaib M. Renal volumes measured by 3-dimensional sonography in healthy fetuses from 20 to 40 weeks. *J Ultrasound Med* 2013;32(3):421-427.
  52. Chang CH, Yu CH, Chang FM, Ko HC, Chen HY. Assessment of fetal adrenal gland volume using three-dimensional ultrasound. *Ultrasound Med Biol* 2002;28(11-12):1383-1387.
  53. Buhimschi CS, Turan OM, Funai EF, Azpurua H, Bahtiyar MO, Turan S, Zhao G, Dulay A, Bhandari V, Copel JA, et al. Fetal adrenal gland volume and cortisol/dehydroepiandrosterone sulfate ratio in inflammation-associated preterm birth. *Obstet Gynecol* 2008;111(3):715-722.
  54. Gindes L, Pretorius DH, Romine LE, Kfir M, D'Agostini D, Hull A, Achiron R. Three-dimensional ultrasonographic depiction of fetal abdominal blood vessels. *J Ultrasound Med* 2009;28(8):977-988.
  55. Turan OM, Turan S, Buhimschi IA, Funai EF, Campbell KH, Bahtiyar OM, Harman CR, Copel JA, Baschat AA, Buhimschi CS. Comparative analysis of 2D versus 3D ultrasound estimation of the fetal adrenal gland volume and prediction of preterm birth. *Am J Perinatol* 2012;29(9):673-680.
  56. Chang CH, Yu CH, Chang FM, Ko HC, Chen HY. Assessment of fetal adrenal gland volume using three-dimensional ultrasound. *Ultrasound in Med Bio* 2002;28(11-12):1383-1387.
  57. Hoffman Sage Y, Lee L, Thomas AM, Benson CB, Shipp TD. Fetal adrenal gland volume and preterm birth: a prospective third-trimester screening evaluation. *J Maternal-Fetal and Neonatal Med* 2015:1-4.
  58. Roelfsema NM, Hop WC, Boito SM, Wladimiroff JW. Three-dimensional sonographic measurement of normal fetal brain volume during the second half of pregnancy. *Am J Obstet Gynecol* 2004;190(1):275-280.
  59. Chang CH, Yu CH, Chang FM, Ko HC, Chen HY. The assessment of normal fetal brain volume by 3-D ultrasound. *Ultrasound Med Biol* 2003;29(9):1267-1272.
  60. Bornstein E, Monteagudo A, Santos R, Keeler SM, Timor-Tritsch IE. A systematic technique using 3-dimensional ultrasound provides a simple and reproducible mode to evaluate the corpus callosum. *Am J Obstet Gynecol* 2010;202(2):201e1-5.
  61. Bornstein E, Monteagudo A, Santos R, Strock I, Tsymbal T, Lenchner E, Timor-Tritsch IE. Basic as well as detailed neurosonograms can be performed by offline analysis of three-dimensional fetal brain volumes. *Ultrasound Obstet Gynecol* 2010;36(1):20-25.
  62. Gindes L, Weissmann-Brenner A, Weisz B, Zajicek M, Geffen KT, Achiron R. Identification of the fetal hippocampus and fornix and role of 3-dimensional sonography. *J Ultrasound Med* 2011;30(12):1613-1618.
  63. Rizzo G, Pietrolucci ME, Capponi A, Arduini D. Assessment of corpus callosum biometric measurements at 18 to 32 weeks gestation by 3-dimensional sonography. *J Ultrasound Med* 2011;30(1):47-53.
  64. Scheier M, Lachmann R, Petros M, Nicolaides KH. Three-dimensional sonography of the posterior fossa in fetuses with open spina bifida at 11-13 weeks gestation. *Ultrasound Obstetrics Gynecol* 2011;38(6):625-629.
  65. Rizzo G, Pietrolucci ME, Mammarella S, Dijmeli E, Bosi C, Arduini D. Assessment of cerebellar vermis biometry at 18 to 32 weeks of gestation by three-dimensional ultrasound examination. *J Maternal-Fetal and Neonatal Med* 2012;25(5):519-522.
  66. Scott JA, Hamzelou KS, Rajagopalan V, Habas PA, Kim K, Barkovich AJ, Glenn OA, Studholme C. 3D morphometric analysis of human fetal cerebellar development. *Cerebellum* 2012;11(3):761-770.
  67. Zeng S, Zhou QC, Zhou JW, Li M, Long C, Peng QH. Volume of intracranial structures on three-dimensional ultrasound in fetuses with congenital heart disease. *Ultrasound Obstetrics Gynecol* 2015;46(2):174-181.
  68. Monteagudo A, Timor-Tritsch IE, Mayberry P. Three-dimensional transvaginal neurosonography of the fetal brain: navigating in the volume scan. *Ultrasound Obstetrics Gynecol* 2000;16(4):307-313.
  69. Kuklisova-Murgasova M, Cifor A, Napolitano R, Papa-georgiou A, Quaghebeur G, Rutherford MA, Hajnal JV, Noble JA, Schnabel JA. Registration of 3D fetal neurosonography and MRI. *Medical Image Analysis* 2013;17(8):1137-1150.